

***Euclid*: Early Release Observations – A preview of the *Euclid* era through a galaxy cluster magnifying lens[★]**

H. Atek^{★1}, R. Gavazzi^{2,1}, J. R. Weaver³, J. M. Diego⁴, T. Schrabback⁵, N. A. Hatch⁶, N. Aghanim⁷, H. Dole⁷, W. G. Hartley⁸, S. Taamoli⁹, G. Congedo¹⁰, Y. Jimenez-Teja^{11,12}, J.-C. Cuillandre¹³, E. Bañados¹⁴, S. Belladitta^{14,15}, R. A. A. Bowler¹⁶, M. Franco¹⁷, M. Jauzac^{18,19,20,21}, G. Mahler^{22,18,19}, J. Richard²³, P.-F. Rocci⁷, S. Serjeant²⁴, S. Toft^{25,26}, D. Abriola²⁷, P. Bergamini^{27,15}, A. Biviano^{28,29}, P. Dimauro^{30,12}, M. Ezziati², J. B. Golden-Marx⁶, C. Grillo^{27,31}, A. C. N. Hughes³², Y. Kang⁸, J.-P. Kneib³³, M. Lombardi²⁷, G. A. Mamon^{1,34}, C. J. R. McPartland^{35,26}, M. Meneghetti^{15,36}, H. Miyatake^{37,38,39}, M. Montes^{40,41}, D. J. Mortlock^{32,42}, P. A. Oesch^{8,26,25}, N. Okabe^{43,44,45}, P. Rosati^{46,15}, A. N. Taylor¹⁰, F. Tarsitano⁸, J. Weller^{47,48}, M. Kluge⁴⁸, R. Laureijs⁴⁹, S. Paltani⁸, T. Saifollahi^{50,51}, M. Schirmer¹⁴, C. Stone⁵², A. Mora⁵³, B. Altieri⁵⁴, A. Amara⁵⁵, S. Andreon⁵⁶, N. Auricchio¹⁵, M. Baldi^{57,15,36}, A. Balestra⁵⁸, S. Bardelli¹⁵, A. Basset⁵⁹, R. Bender^{48,47}, C. Bodendorf⁴⁸, D. Bonino⁶⁰, E. Branchini^{61,62,56}, M. Brescia^{63,64,65}, J. Brinchmann^{66,67}, S. Camera^{68,69,60}, G. P. Candini⁷⁰, V. Capobianco⁶⁰, C. Carbone³¹, V. F. Cardone^{30,71}, J. Carretero^{72,73}, S. Casas⁷⁴, F. J. Castander^{75,76}, M. Castellano³⁰, S. Cavuoti^{64,65}, A. Cimatti⁷⁷, C. J. Conselice¹⁶, L. Conversi^{78,54}, Y. Copin⁷⁹, L. Corcione⁶⁰, F. Courbin³³, H. M. Courtois⁸⁰, A. Da Silva^{81,82}, H. Degaudenzi⁸, A. M. Di Giorgio⁸³, J. Dinis^{81,82}, M. Douspis⁷, F. Dubath⁸, X. Dupac⁵⁴, S. Dusini⁸⁴, A. Ealet⁷⁹, M. Farina⁸³, S. Farrens¹³, S. Ferriol⁷⁹, S. Fotopoulou⁸⁵, M. Frailis²⁸, E. Franceschi¹⁵, S. Galeotta²⁸, W. Gillard⁸⁶, B. Gillis¹⁰, C. Giocoli^{15,87}, P. Gómez-Alvarez^{88,54}, A. Grazian⁵⁸, F. Grupp^{48,47}, L. Guzzo^{27,56}, M. Hailey⁷⁰, S. V. H. Haugan⁸⁹, J. Hoar⁵⁴, H. Hoekstra⁹⁰, M. S. Holliman¹⁰, W. Holmes⁹¹, I. Hook⁹², F. Hormuth⁹³, A. Hornstrup^{94,35}, P. Hudelot¹, K. Jahnke¹⁴, M. Jhabvala⁹⁵, E. Keihänen⁹⁶, S. Kermiche⁸⁶, A. Kiessling⁹¹, T. Kitching⁷⁰, R. Kohley⁵⁴, B. Kubik⁷⁹, K. Kuijken⁹⁰, M. Kümmel⁴⁷, M. Kunz⁹⁷, H. Kurki-Suonio^{98,99}, O. Lahav¹⁰⁰, D. Le Mignant², S. Ligori⁶⁰, P. B. Lilje⁸⁹, V. Lindholm^{98,99}, I. Lloro¹⁰¹, D. Maino^{27,31,102}, E. Maiorano¹⁵, O. Mansutti²⁸, O. Marggraf¹⁰³, K. Markovic⁹¹, N. Martinet², F. Marulli^{104,15,36}, R. Massey^{18,19}, S. Maurogordato¹⁰⁵, H. J. McCracken¹, S. Mei¹⁰⁶, Y. Mellier^{34,1}, E. Merlin³⁰, G. Meylan³³, M. Moresco^{104,15}, L. Moscardini^{104,15,36}, R. Nakajima¹⁰³, R. C. Nichol⁵⁵, S.-M. Niemi⁴⁹, C. Padilla¹⁰⁷, K. Paech⁴⁸, F. Pasian²⁸, J. A. Peacock¹⁰, K. Pedersen¹⁰⁸, W. J. Percival^{109,110,111}, V. Pettorino⁴⁹, S. Pires¹³, G. Polenta¹¹², M. Poncet⁵⁹, L. A. Popa¹¹³, L. Pozzetti¹⁵, F. Raison⁴⁸, A. Renzi^{114,84}, J. Rhodes⁹¹, G. Riccio⁶⁴, E. Romelli²⁸, M. Roncarelli¹⁵, R. Saglia^{47,48}, D. Sapone¹¹⁵, P. Schneider¹⁰³, A. Secroun⁸⁶, G. Seidel¹⁴, S. Serrano^{76,116,75}, C. Sirignano^{114,84}, G. Sirri³⁶, J. Skottfelt¹¹⁷, L. Stanco⁸⁴, P. Tallada-Crespi^{72,73}, H. I. Teplitz¹¹⁸, I. Tereno^{81,119}, R. Toledo-Moreo¹²⁰, I. Tutusaus¹²¹, L. Valenziano^{15,122}, T. Vassallo^{47,28}, G. Verdoes Kleijn⁵¹, A. Veropalumbo^{56,62,123}, Y. Wang¹¹⁸, E. Zucca¹⁵, C. Baccigalupi^{29,28,124,125}, C. Burigana^{126,122}, G. Castignani¹⁵, Z. Sakr^{127,121,128}, V. Scottez^{34,129}, M. Viel^{29,28,125,124,130}, P. Simon¹⁰³, D. Stern⁹¹, J. Martín-Fleitas⁵³, and D. Scott¹³¹

(Affiliations can be found after the references)

May 23, 2024

ABSTRACT

We present the first analysis of the *Euclid* Early Release Observations (ERO) program that targets fields around two lensing clusters, Abell 2390 and Abell 2764. We use VIS and NISP imaging to produce photometric catalogs for a total of $\sim 500\,000$ objects. The imaging data reach a 5 σ typical depth in the range 25.1–25.4 AB in the NISP bands, and 27.1–27.3 AB in the VIS band. Using the Lyman-break method in combination with photometric redshifts, we search for high-redshift galaxies. We identify 30 Lyman-break galaxy (LBG) candidates at $z > 6$ and 139 extremely red sources (ERSs), most likely at lower redshift. The deeper VIS imaging compared to NISP means we can routinely identify high-redshift Lyman breaks of the order of 3 magnitudes, which reduces contamination by brown dwarf stars and low-redshift galaxies. The difficulty of spatially resolving most of these sources in 0.3 pix⁻¹ imaging makes the distinction between galaxies and quasars more challenging. Spectroscopic follow-up campaigns of such bright sources will help constrain both the bright end of the ultraviolet galaxy luminosity function and the quasar luminosity function at $z > 6$, and constrain the physical nature of these objects. Additionally, we have performed a combined strong lensing and weak lensing analysis of A2390, and demonstrate how *Euclid* will contribute to better constraining the virial mass of galaxy clusters. From these data, we also identify optical and near-infrared counterparts of known $z > 0.6$ clusters, which exhibit strong lensing features, establishing the ability of *Euclid* to characterize high-redshift clusters. Finally, we provide a glimpse of *Euclid*'s ability to map the intracluster light out to larger radii than current facilities, enabling a better understanding of the cluster assembly history and mapping of the dark matter distribution. This initial dataset illustrates the diverse spectrum of legacy science that will be enabled by the *Euclid* survey.

Key words. Galaxies: clusters: general; Galaxies: high-redshift; Gravitational lensing: strong; Gravitational lensing: weak; reionization; Cosmology: observations

1. Introduction

Beyond its primary cosmology science goals, the *Euclid* mission will provide a sample of more than 200 000 high-redshift galaxies across 14 000 deg² of extragalactic sky (Euclid Collaboration: Scaramella et al. 2022; Euclid Collaboration: Mellier et al. 2024). This will enable a wide array of legacy science. In particular, the near-infrared (NIR) capabilities of the NISP (Near-Infrared Spectrometer and Photometer) instrument (Euclid Collaboration: Jahnke et al. 2024) and the wide optical coverage of the visible instrument (VIS; Euclid Collaboration: Cropper et al. 2024), allow us to detect star-forming galaxies and quasars during the epoch of reionization, ranging from $z = 6$ to $z = 9$.

The Deep Survey component of *Euclid* will map a total of 53 deg², namely in the Euclid Deep Field North (EDF-N), the Euclid Deep Field South (EDF-S), and the Euclid Deep Field Fornax (EDF-F), down to a 5σ limiting magnitude of ~ 26.2 in the NIR and 28.5 in the visible. In comparison, the widest areas covered by the *Hubble* Space Telescope (HST) and the JWST, are of the order of 1 deg². The Wide Survey will reach a 5σ magnitude depth of 26.2 in VIS/ I_e and 24.5 in all NISP bands. The sky coverage is optimized to have maximum overlap with ground-based ancillary data, which greatly help with photometric redshift estimates (Euclid Collaboration: van Mierlo et al. 2022; Euclid Collaboration: Scaramella et al. 2022; Euclid Collaboration: Mellier et al. 2024).

One of the main legacy science components is the identification of distant galaxies and quasars at redshifts greater than 6. This will rely primarily on the detection of Lyman break in the VIS I_e filter at $z \sim 7$, and in one of the NISP filters at higher redshifts. Based on the current survey design, more than 10 000 galaxies are expected at $z \sim 7$ and up to 2000 galaxies at $z \sim 8$, assuming a Schechter UV luminosity function (Bouwens et al. 2015). Over 100 quasars are expected over $7.0 < z < 7.5$, and about 25 quasars beyond $z = 7.5$, assuming a declining quasar luminosity function beyond $z = 6$ (Euclid Collaboration: Barnett et al. 2019). The brightness of these sources, $J_e \gtrsim 21$ opens up spectroscopic follow-up opportunities with ground-based telescopes to robustly determine the bright-end of the galaxy and the quasar luminosity functions. In addition, *Euclid* will observe hundreds of lensing clusters, enabling the detection of magnified galaxies and quasars at $z > 6$.

Here we present an overview and early results of the Early Release Observations (ERO) program Magnifying Lens, which targets two cluster fields, Abell 2390 and Abell 2764 (hereafter A2390 and A2764, respectively). This program represents a unique opportunity to showcase the science questions that *Euclid*'s observations of galaxy clusters will be able to address. This paper serves as an introduction to these initial measurements and findings in both cluster and blank fields. Specifically, it illustrates how *Euclid* will identify high-redshift dropout sources in both blank regions and behind lensing clusters. A companion paper is dedicated to the search for high-redshift candidates at $z > 6$ (see Weaver et al. 2024). We also take this opportunity to investigate the challenges in robustly identifying high-redshift objects and the potential sources of contamination. These observations also offer a glimpse into the combination of strong lensing and weak lensing analyses to map the mass distribution in lensing clusters that will be uncovered in *Euclid* data.

Furthermore, *Euclid* offers a wide field of view that is necessary to map the intracluster light in the outer regions of the cluster.

The paper is organized as follows. In Sect. 2, we describe the ERO program and the characteristics of the targets. Section 3 is dedicated to the observations and data reduction of the VIS and NISP imaging. The photometric catalogs that are released with this publication are described in Sect. 4. In terms of scientific exploitation of this dataset, we present the VIS-dropout sources in Sect. 5, the weak lensing analysis of A2390 in Sect. 6 and a combined strong lensing and weak lensing analysis of A2390 in Sect. 7. The search for high-redshift galaxy clusters is discussed in Sect. 8, while the study of the intracluster light in A2390 is presented in Sect. 9. A summary of this work is presented in Sect. 10. We assume a flat Λ CDM cosmology with $H_0 = 70$ km s⁻¹ Mpc⁻¹, $\Omega_m = 0.3$ and $\Omega_\Lambda = 0.7$. All magnitudes are in the AB system.

2. The ERO program

The ERO program was designed to obtain a comprehensive dataset spanning across multiple interdisciplinary working groups, including, but not limited to, those focused on the primordial universe, galaxy evolution and active galactic nuclei (AGN), strong and weak lensing, and galaxy cluster physics. These observations provide a first glimpse of the *Euclid* imaging data, with the aim of facilitating the inaugural evaluation of tools, data quality, and methodologies developed over the preceding decade. The early investigation of the dataset will help reveal and address challenges and limitations for a full scientific exploitation of the prime mission survey data.

The ERO observations for this project were taken on 28 November 2023, as part of the performance-verification (PV) phase of the *Euclid* mission.

2.1. Abell 2390

The first ERO target is the galaxy cluster A2390, RA=21^h 53^m 35^s.50, Dec=17° 41' 41".54, located at a redshift of $z = 0.228$ (Sohn et al. 2020). It is one of the richest clusters in the Abell catalog (Pello et al. 1991; Le Borgne et al. 1991; Feix et al. 2010) with an estimated mass M_{200}^1 ranging from $1.53 \times 10^{15} M_\odot$ from weak lensing (Okabe & Smith 2016) to $1.84 \times 10^{15} M_\odot$ from the projected phase-space of galaxies (Sohn et al. 2020). It includes multiple well-studied lensing arcs (e.g., Olmstead et al. 2014; Richard et al. 2021). It also displays bright X-ray emission with a luminosity of 2.7×10^{45} erg s⁻¹ in the 0.1–2.4 keV range (Boehringer et al. 1998). The *Euclid* pointing is centered on the cluster, which represents a small fraction of the total area (cf. Fig. 1).

2.2. Abell 2764

The second ERO target is a random field on the sky that happens to cover the galaxy cluster A2764 in a corner of the field (see Fig. 2). The coordinates of the center of the field are RA=0^h 22^m 50^s.

¹ M_{200} is the mass within the virial radius, which is where the mass density of the system is 200 times the critical density of the Universe.

* This paper is published on behalf of the Euclid Consortium

** e-mail: atek@iap.fr




Fig. 1. *Euclid* color composite image of the A2390 field. The blue channel is the I_E image; green is Y_E , and red is H_E . The whole field covers about 0.5 deg^2 . The inset is centered on A2390 and covers $2' \times 2'$, which is approximately the field of view of the Wide Field Camera 3 (WFC3) NIR channel onboard HST. The whole *Euclid* field identifies approximately 250 000 sources in the CHL_MEAN detection image based on the Y_E , J_E , and H_E coadd mosaic.

1347, Dec= $-49^\circ 15' 59''.87$. The galaxy cluster was first identified by Abell et al. (1989). It is located at a redshift of $z = 0.07$ (Katgert et al. 1996), and is less massive ($M_{500} = 9.2 \times 10^{13} M_\odot$) than A2390 (Serenio & Ettori 2017). It is also far less studied and characterized than A2390. Additional clusters can be identified in the field and are described in Sect. 8.

3. Observations and data reduction

3.1. *Euclid* observations

For each of the Abell clusters, we obtained three reference observing sequences (ROS; *Euclid* Collaboration: Scaramella et al. 2022), implying 3 times the nominal *Euclid* Wide Survey exposure time. The duration of each ROS is about 70.2 min. A standard 4-point dither pattern, with an offset of $120'' \times 220''$, was applied for each ROS. In addition, we adopted a $3' \times 3'$ dither between the individual ROS to maximize the depth. The dithered pointings can be seen in the weight maps of the NISP imaging




Fig. 2. Color-composite image of the A2764 field using the I_E image for the blue channel, Y_E for green, and H_E for red. The image field of view is about 0.5 deg^2 and contains nearly 250 000 objects based on the NISP coadd detection image. The inset shows a closeup of edge-on and face-on galaxies in the A2764 cluster.

presented in Fig. 3. The large dithers fill the detector gaps and allow us to obtain continuous coverage of the field. The field of view of each combined mosaic is about 0.75 deg^2 .

The data reduction process is described in detail in [Cuillandre et al. \(2024\)](#). Individual frames were stacked following two different approaches. We produced compact-sources stacks with the *AstrOmatic SWarp* software ([Bertin et al. 2002](#)) using the individual multiple extension fits (MEF) files and the associated weight maps. The procedure achieves an astrometric accuracy of 5.1/15.2 milliarcseconds (mas) in I_E/H_E , respectively, for A2390 and 7.9/15.8 mas for A2764. A mesh size of 64 pixels (with a smoothing factor of 3) was adopted to subtract the background

in input images that feature large extended low-surface brightness (LSB) emission. The second flavor of the data reduction also uses *SWarp* on the individual frames and does not perform any internal background subtraction, thereby conserving LSB features such as interstellar cirrus emission. This is particularly useful for studies of diffuse intracluster light. The sigma-clipping algorithm used to build these extended-emission stacks slightly affects the photometry of very compact sources, such as stars, causing an additional 5% photometric error in this case. The images were masked to contain only overlapping regions in the dithered frames of both VIS and NISP. The final sampling of VIS and NISP images is $0''.1 \text{ pix}^{-1}$ and $0''.3 \text{ pix}^{-1}$, respectively.




Fig. 3. Weight maps of the *Euclid* observations of the lensing clusters A2390 (left) and A2764 (right). These examples are associated with the J_E observations. The maps demonstrate the large dither pattern adopted for these ERO observations and the sky coverage of each field.

A composite color image of A2390 is shown in Fig. 1. One can easily appreciate the gain in area, with one pointing only, over the limited survey capabilities of HST. The zoom-in inset illustrates the HST/WFC3 field of view over the center of the field (the lensing cluster). The total single pointing area of 0.75 deg^2 (cropped to 0.5 deg^2 in the color image) is larger than that of the entire Cosmic Assembly Near-infrared Deep Extragalactic Legacy Survey (CANDELS) area (Koekemoer et al. 2011), one of the largest HST extragalactic legacy programs. We also show a composite color image of A2764 in Fig. 2.

3.2. Ancillary observations

These two ERO fields benefit from various space-based and ground-based ancillary data. For A2390, there is 6-band (B, V, R_c, i, I_c, z') imaging data (Miyazaki et al. 2002) obtained with the Suprime-Cam instrument on the 8.2-m Subaru telescope. The data, which were reduced by the processing pipeline SDFRED (Yagi et al. 2002; Ouchi et al. 2004), cover a $28' \times 34'$ region centered on the brightest cluster galaxy (BCG, Okabe & Smith 2016). In addition, we also consider Canada France Hawaii Telescope (CFHT) WIRCam J -band observations, obtained in April-May-September 2010 covering the inner $24' \times 24'$ region, and a CFHT Megacam u -band (first generation MP9301) image taken in August 2007 for a total integration time of 2000 s and covering the whole ERO field of view. Details of the reduction, astrometric and photometric calibration of these ancillary data will be presented in a subsequent paper dedicated to weak lensing and photometric redshifts.

Previously, the cluster was imaged with HST using both the Wide Field Planetary Camera 2 (WFPC2) in F555W and F814W, and the Advanced Camera for Surveys (ACS) in F850LP (see Ota et al. 2012). There are also VLT/MUSE integral-field spectroscopy (PID: 094.A-0115), which support the reported strong lensing mass modelling presented in Richard et al. (2021).

In addition, the cluster core of A2390 has been observed with ALMA Bands 3-5 (PID 2021.1.00766.S) and SITELLE on the CFHT (see Liu et al. 2021, and references therein), which reveal

Table 1. Limiting magnitudes at 5σ reached in each *Euclid* filter of the A2390 and A2764 fields. The depth measurements assume an aperture diameter of $0''.6$ and $0''.3$ for NISP and VIS, respectively.

Field	RA [deg]	Dec [deg]	I_E	Y_E	J_E	H_E
A2390	328.397	+17.709	27.01	25.18	25.22	25.12
A2764	5.713	-49.249	27.26	25.30	25.41	25.21

an $H\alpha$ emission line in a cluster member galaxy. Furthermore, the cluster core of A2390 has been mapped with the SCUBA-2 submillimeter camera by Cowie et al. (2022) to 1σ depths of 0.5 mJy at $850 \mu\text{m}$ and $2.5\text{--}3.5 \text{ mJy}$ at $450 \mu\text{m}$. Of the 59 submillimeter galaxies in the field, 44 have a plausible cross-identification (within $2''$) in the ERO data.

Both fields have been observed with the Dark Energy Camera (DECam; Flaugher et al. 2015) on the Blanco telescope at the Cerro Tololo Inter-American Observatory. A2764 lies within the Dark Energy Survey (DES; Flaugher 2005; Dark Energy Survey Collaboration et al. 2016) footprint, with typical 10σ depths ($1''.95$ aperture) for the $griz$ -bands in DR2² of 24.7, 24.4, 23.8 and 23.1, respectively (Abbott et al. 2021). Typical image qualities for these DES bands are $1''.11$, $0''.95$, $0''.88$ and $0''.83$. DES data will be used to partner the Southern regions of the *Euclid* Wide Survey during *Euclid*'s first data release, and so this largely-blank field provides an early example of those forthcoming combined data. Meanwhile, A2390 has been observed in grz -bands as part of the DESI Legacy Survey (DLS;³ Dey et al. 2019). Typical depths (5σ , extended source) and image qualities of the DLS images are, 23.72, 23.27, 22.22 and $1''.29$, $1''.18$, $1''.11$, in the grz -bands, respectively.




Fig. 4. Photometric errors as a function of the magnitude for all sources in the catalog for A2390 (left) and A2764 (right). Each curve represents the median error for each VIS and NISP band, while the corresponding shaded area is the 1- σ dispersion. The error measurements are described in Sect. 4.1.

4. Photometric catalogs

4.1. Point-source aperture photometry

The combination of NISP NIR bands (Y_E , J_E , and H_E) and ultra-deep VIS optical imaging in I_E , about 2 magnitudes deeper, enables immediate searches for ultra-red objects with no appreciable optical flux. Such objects include high-redshift galaxies and quasars ($z \gtrsim 6$) as well as dusty and/or quiescent galaxies, cool stars and brown dwarfs at lower redshifts. The former were a key scientific goal of this ERO program. To enable rapid searches for such objects (see [Weaver et al. 2024](#)), we first co-add the compact-source version of $Y_E + J_E + H_E$ mosaics to produce a CHI_MEAN detection image using SWarp ([Bertin et al. 2002](#)). From that image, we detect 501 994 objects (249 319 in A2764 and 252 675 in A2390) using Pythonic SourceExtractor ([Bertin & Arnouts 1996](#); [Barbary 2016](#)) assuming a threshold of 1.5σ within at least three contiguous pixels of the detection image smoothed by a Gaussian kernel with a $0''.45$ (1.5 pixels) FWHM.⁴ De-blending is configured with 2×10^5 thresholds requiring a contrast of 10^{-5} . No cleaning is applied as it is liable to remove real sources of interest. Photometry is extracted at the source positions by summation of pixels within several circular apertures with diameters: $0''.1$, $0''.2$, $0''.3$, $0''.6$, $1''.0$, $1''.2$, $1''.5$, and $2''.0$. Uncertainties are estimated strictly from extractions on the weight map, whose distribution in effective per-pixel error matches well to the observed noise level in the images. Given the relative similarity of NISP PSFs measured by stacking point sources in the images, we elect to simply correct all apertures to total flux, assuming point-like light distributions outside the apertures. Although we provide similarly corrected total photometry for VIS, it is the signal-to-noise ratio that is most important to our objective, where aperture corrections are superfluous.

Therefore, given the uncertainty in recovering total fluxes, in particular for extended objects, we strongly caution against combining this photometry with other pre-compiled catalogs as colors are not guaranteed to be robust. Figure 4 shows the typical magnitude uncertainty in each band as a function of magnitude for all the sources detected and included in the photometric catalog. It highlights, in particular, the stable photometry across a wide flux range and the significant difference in depth between the VIS and the NISP observations. Magnitude zero point uncertainties are estimated to be below a 10% level in these data. We estimated the limiting magnitude for each of the *Euclid* bands of our observations by measuring the flux in 10 000 empty apertures randomly distributed in the A2390 and A2764 images. The standard deviation of the flux distribution provides the typical depths quoted in Tab. 1.

We provide a $I_E, Y_E, J_E,$ and H_E photometric catalog of the two ERO fields, which includes unique and field-specific IDs, coordinates, basic shape parameters, aperture and total fluxes, aperture-to-total corrections, and SourceExtractor-like flags. All photometric measurements are corrected for Galactic extinction using the extinction curves of [Fitzpatrick & Massa \(2007\)](#). A total of 73 841 unreliable sources (14.7% of the total) likely to be artifacts near mosaic edges (where only a single dither contributes to the mosaic) or near bright stars are flagged by `use_phot=0`. See the accompanying README file for details of the source detection, photometry, and catalog contents. A case study utilizing this catalog to identify NISP-only sources is presented in [Weaver et al. \(2024\)](#).

4.2. SourceXtractor++

We separately derive photometric catalogs containing morphological parameters based on the recent re-implementation of SExtractor ([Bertin & Arnouts 1996](#)), called SourceXtractor++ (hereafter SE++; [Kümmel et al. 2022](#); [Bertin et al. 2022](#)). Among several other features, this code allows the user to fit PSF-convolved parametric models

² <https://des.ncsa.illinois.edu/releases/dr2>

³ <http://legacysurvey.org/>

⁴ A discussion on thresholds and kernel sizes will be presented in Zalesky et al. in prep.

describing the surface brightness profile of every single source in the images, the fitting being performed simultaneously on several exposures grouped into a number of photometric filters or bands. The PSF, and its spatial variations across the field of view, are described with the PSFEX tool (Bertin 2011). In the framework of the Euclid Morphology Challenges (Euclid Collaboration: Merlin et al. 2023; Euclid Collaboration: Bretonnière et al. 2023), we have shown that SE++ can recover accurate multi-band photometry and morphological parameters, such as half-light radius, ellipticity, orientation, and Sérsic index. Mandelbaum et al. (2015) and Euclid Collaboration: Martinet et al. (2019) have previously shown that recovered shapes obtained with the model-fitting engine of SExtractor meet the weak lensing standards (with a proper weighting of source ellipticities) that is sufficient for the scopes of cluster analysis and stage III weak lensing surveys. A forthcoming weak lensing study of A2390 will investigate in detail the merits of weak lensing shape derivations with three methods: SE++, LENS MC, currently the main cosmic shear measurement adopted for the first data release in the OU-SHE part of the *Euclid* Ground Segment (see Congedo et al. 2024, in prep.), and KSB++ (Kaiser et al. 1995; Luppino & Kaiser 1997; Hoekstra et al. 1998).

We ran SE++ in two settings, but source detection was always performed in the VIS I_E band. In the first run, only the VIS band was used to constrain a single Sérsic profile. This provides shapes that can readily be used for weak lensing, and best-fit sizes provide a complementary star/galaxy discriminator. Then, a two-component bulge+disk (i.e. de Vaucouleurs + exponential) concentric model was also fit to all the ground-based, VIS and NISP bands at once, i.e. $u, B, V, R_c, i, I_c, z, I_E, Y_E, J_E, H_E$. The orientation of the disk and the bulge was forced to coincide, but their axis ratios were left free. The bulge and disk half-light radii were constrained to be constant across wavelengths, but the bulge-to-total flux ratio was allowed to vary. Furthermore, in order to account for the long time span between SuprimeCam (early 2000') and *Euclid* (November 2023) observations, we allow for proper motions for all sources.

Both runs rely on the same PSF determination based on PSFEX. In addition, further measurements were performed besides those that depend on model fitting. In particular, aperture photometry in 1", 3", and 6" diameter apertures was computed, as well as source centroid and isophotal moments, in line with standard SExtractor measurements.

We applied a careful masking of sources in the vicinity of bright stars with an automated scaling with stellar magnitude of the size of polygons, aimed to mask out the extended diffraction spikes (Fig. 5). A model of ghosts due to reflections onto the dichroic, which splits light between the VIS and NISP instruments, was also defined. Ghosts from Gaia-DR3 stars brighter than $G_{RP} = 12$ are automatically flagged as a disk of 8" radius, shifted by about 1' with respect to the source star. Likewise, stars (core and spikes) between Gaia magnitudes $G_{RP} = 8.5$ and $G_{RP} = 18.5$ are automatically masked. Since the brightness profile along diffraction spikes drops with the angular distance to the star center θ as $\theta^{-2.2}$, we scale the size of corresponding masks as $\exp(-0.41 G_{RP})$. After automated generation of masks, further tuning was done by hand. In the case of A2390, before masking, the total relevant area is 0.75 deg². After masking out those regions, the useful VIS area reduces to 0.71 deg². The unmasked area with deep multi-band SuprimeCam data further reduces to 0.21 deg².

Multi-band photometry at fixed positions allows us to reach greater depth in the filters that are not the VIS I_E detection image. We reach a 0.1 magnitude uncertainty on model magnitudes for




Fig. 5. *Euclid* I_E image (1'5 × 1'0) in the A2390 field of view, showing masked area (inside polygons). The image shows diffraction spikes, saturation bleeds, core region, and two reflection ghosts of 8" radius (in red).

A2390 (total bulge+disk decomposition) at 24.14, 26.19, 25.80, 26.41, 25.48, 25.64, and 24.76 for filters $u, B, V, R_c, i, I_c,$ and z , respectively. This depth, well-matched to *Euclid* depth, was achieved thanks to exquisite observing conditions, during which seeing reached 0'905, 0'736, 0'572, 0'561, 0'516, 0'682, and 0'570, respectively. Once combined with VIS and NISP *Euclid* photometry, this will enable computation of accurate photometric redshifts, as will be shown in the dedicated weak lensing paper. As an illustration, note the tight red sequence of cluster member galaxies in the left panel of Fig. 6. For details, refer to the README file associated with the catalog release.

5. High-redshift dropout sources

The relative depths of VIS and NISP imaging, which together cover the entire 0.55–2 μm spectral range, means we can identify high-redshift ($z > 6$) dropout sources, that are well detected in NISP and remain undetected in the deeper VIS data. Here we present a summary of the results from the selection of high-redshift sources. A detailed analysis of these findings, along with the challenges in identifying high-redshift candidates, is provided in a dedicated paper (Weaver et al. 2024). We achieve a 5σ depth in the range 25.1–25.4 in the NISP bands, and 27.1–27.3 in the VIS image, assuming circular apertures with diameter 0'6 and 0'3, respectively (see Table 1). In addition, we require our sources to be detected at a 3σ level in the Y_E band and to be simultaneously undetected in the I_E band at a 1.5σ level. This leads to a source selection that is based on a minimum Lyman break of approximately 2 magnitudes, which is significantly higher than most Lyman-break selection criteria used in HST, and JWST, identification of $z > 6$ galaxies (e.g., Bouwens et al. 2015; Finkelstein et al. 2015; Atek et al. 2023). A stronger dropout criterion significantly reduces contamination by red low-redshift interlopers. However, ERSs can still enter our selection window but can be differentiated from LBGs by their clearly extreme red colors compared to the relatively flat or blue UV slopes observed in LBGs above the Lyman break (cf. Fig. 7) and the existence of a secondary peak at low redshift in the posterior probability distribution of their photometric redshift (see below). The ERSs are an interesting population that constitutes its own science case at lower redshift. The strong Lyman break does not completely ex-




Fig. 6. *Left:* Galaxy number density in the plane of Subaru SuprimeCam $B - V$ color vs. *Euclid* I_E -band magnitude, from bulge+disk light profile fits with SourceExtractor++ (see text). Red stars indicate spectroscopic cluster members of A2390 determined by Sohn et al. (2020) and MUSE observations of the core of A2390. The dashed region indicates the cluster red sequence based on the spectroscopic members and extrapolated to fainter magnitudes. *Right:* Sky location of galaxies enclosed by the red-sequence selection in the left-hand panel over the joint 0.21 deg^2 area covered by *Euclid* and Subaru SuprimeCam. The size of each point scales inversely with I_E magnitude to indicate the distribution of stellar mass in red galaxies at the cluster redshift.

clude some cool brown dwarf stars, especially given the limited spatial resolution of NISP, which only enables resolving multi-component bright galaxies, which might be common at high redshift (Bowler et al. 2017). To minimize contamination by artifacts, we also require the sources to be compact by imposing a flux ratio between two apertures of $J_E (1''.5)/J_E (0''.6) < 1.3$. We have verified that the criterion does not exclude real sources from the selection. Furthermore, in order to mitigate contamination by persistence at NIR wavelengths from previous exposures, we performed a time series analysis on the individual frames. We removed all sources that showed a typical declining light curve (for more details, see Weaver et al. 2024).

This color selection is complemented with photometric redshift estimates using the spectral energy distribution (SED) fitting tool eazy-py (Brammer et al. 2008, 2022). We assume a redshift range of $0.01 < z < 15$ and a flat luminosity prior using the agn_blue_sfhz set of galaxy templates. All the sources have a posterior distribution function (PDF) that is consistent with the $z > 6$ solution, with no significant secondary solution at low redshift. The full procedure and the dropout sample are described in detail in Weaver et al. (2024). Combining the two ERO fields, we identify a total of 30 LBGs and 139 ERSs, from a total of about 500 000 sources. The absolute magnitudes of the LBGs ranges between $M_{UV} = -21.9$ and $M_{UV} = -23.6$. In Fig. 7, we show an example for each of these categories. The top row shows the photometry in each of the NISP and VIS filters and a composite color image of the Y_E, J_E, H_E images. We also show the best-fit solutions of the photometric data derived with eazy-py, and the associated probability distribution. The LBG candidate has a clear high-redshift solution centered at $z \sim 7.7$, whereas, the ERS has a significant low-redshift solution around $z \sim 1.7$.

These early results, based on a small portion of the sky relative to the full survey, highlight the crucial role that *Euclid* will play in identifying the brightest galaxies during the epoch of reionization. Given the uncertainties on the exact shape of the bright-end of the UV luminosity function at $z \sim 8-10$ (Bowler et al. 2020), on the one hand, and the apparent excess of UV-

bright galaxies at $z > 10$ (Harikane et al. 2023; Chemerynska et al. 2023; Adams et al. 2023; Donnan et al. 2024), on the other hand, *Euclid* imaging and subsequent spectroscopic follow-up will be instrumental in constraining the exact abundance of bright high-redshift galaxies, and the exact shape of the bright end of the UV luminosity function at $z = 6-9$. The cross-correlation with $\text{Ly}\alpha$ emitters will allow the identification of giant ionized bubbles, of the order of 10 Mpc in size, and map the ionization state of the IGM. In addition to bright LBGs, the *Euclid* data will also allow the identification of bright quasars in the deep and wide surveys. The ability of current data to distinguish between high-redshift quasars and galaxies remains limited in the absence of follow-up spectroscopy. *Euclid* grism spectroscopy will be limited to sources brighter than 22 mag. These limitations also apply to potential contaminating sources at low redshifts, such as cool brown dwarf stars (Weaver et al. 2024). Again, the relative depth of VIS imaging compared to NISP, which leads to very strong breaks between these two wavelengths, reduces the contamination rate of Galactic dwarfs, but will remain an important source of contamination (Euclid Collaboration: Barnett et al. 2019). Fortunately, the brightness of these sources makes them ideal candidates for follow-up spectroscopy from the ground in order to confirm their redshifts. Determining the number, density, and luminosity of these sources will motivate follow-up campaigns, with JWST for example, to constrain the nature of the brightest sources in the early Universe, including galaxies and quasars.

6. Weak lensing analysis

Weak gravitational lensing (WL) is one of *Euclid*'s primary cosmological probes. By measuring the shapes of background galaxies, WL analyses allow for the reconstruction of the projected foreground mass distribution (e.g. Bartelmann & Schneider 2001), which is dominated by invisible dark matter. The WL signature of one of the targets of this ERO program, A2390, has previously been studied using ground-based WL data (e.g. von der Linden et al. 2014; Applegate et al. 2014; Okabe &




Fig. 7. The identification of VIS-dropout candidates in the A2390 and A2764 fields. The **top panel** shows an example of an LBG candidate at a redshift of $z > 6$, while the **bottom panel** shows an ERS, most likely located at a lower redshift. For each of these sources, the top row shows the cutout images of the source in each VIS and NISP filters, together with a NISP stack and a color-composite image based on the Y_E , J_E , and H_E filters. The size of these images is about $7'' \times 7''$. The bottom figure illustrates the best-fit model (blue and orange curves) to the *Euclid* photometry (colored circles with associated 1σ uncertainties). The gray horizontal bars denote the 3σ depth of each band, while the down-pointing triangle marks the upper limit of the I_E band. The left inset presents the posterior distribution function for the photometric redshift (see Weaver et al. 2024, for details).

Smith 2016). However, given their outstanding depth and resolution, the new *Euclid* observations of the cluster provide an excellent opportunity to showcase *Euclid*'s WL capabilities. For this, we have measured WL galaxy shapes in the A2390 VIS observations using different shape measurement algorithms in-

cluding LENS_{MC} (which is designated as the primary shape measurement of the *Euclid* Data Release 1), SE++ (see Sect. 4.2), and KSB+ (Kaiser et al. 1995; Luppino & Kaiser 1997; Hoekstra et al. 1998). The details of this analysis will be presented in a separate paper. Here we briefly highlight first results from

the KSB+ analysis, which employs the Erben et al. (2001) implementation of the KSB+ algorithm, including the shear bias calibration from Hernández-Martín et al. (2020), as used, e.g., for the analysis of HST WL observations of distant galaxy clusters from the South Pole Telescope in Schrabback et al. (2018, 2021) and Zohren et al. (2022). After applying KSB+ shape measurement selections, removing masked objects in the vicinity of bright stars (see Sect. 4.2) and very extended galaxies, and requiring a flux signal-to-noise ratio $S/N > 10$ (defined as the FLUX_AUTO/FLUXERR_AUTO ratio from SExtractor), as well as a magnitude in the range $22 < I_e < 26.5$, the resulting catalog provides a WL density of 56.5 sources per arcmin² when averaged within the central 33.3×33.3 region around the cluster center. As an initial result of this analysis, we present a reconstruction of the convergence field κ in Fig. 8. The convergence maps the foreground mass distribution, but is scaled according to the geometric lensing efficiency (e.g. Bartelmann & Schneider 2001). In particular, we apply a Wiener-filtered reconstruction following the procedures described in McInnes et al. (2009), Simon et al. (2009), and Schrabback et al. (2018). Here, the signature of the cluster is detected with high significance in the curl-free E-mode reconstruction (left panel of Fig. 8). In the right panel of Fig. 8, we additionally show a B-mode reconstruction derived from the curl component of the WL shear field. To first order, gravitational lensing only creates curl-free E modes. Accordingly, the B-mode reconstruction provides an assessment for the level of noise (and potential residual systematics) in the reconstruction.

7. Weak plus strong lensing analysis

Euclid WL data can be combined with existing strong lensing (SL) constraints (or new multiple-image systems identified by *Euclid*) enabling a joint SL+WL analysis. SL constraints are useful at constraining the inner region of clusters (typically the central 300 kpc), but do not constrain the outer regions of clusters, thus limiting our ability to estimate their virial masses. *Euclid*'s WL measurements can extend the analysis of the lensing signal to beyond the virial radius. When several multiply-imaged galaxies exist at different redshifts, the combination of both data sets breaks the mass-sheet degeneracy (e.g. Schneider & Er 2008). Among the ERO targets, A2390 offers a unique opportunity to test the capacity of *Euclid* at measuring the virial mass of a massive cluster. This cluster contains 11 multiply-imaged galaxies with confirmed spectroscopic redshifts (Richard et al. 2021). These galaxies produce 23 multiple images that result in 46 SL positional constraints. We combine these constraints with WL measurements from *Euclid*. We rebin the WL data in 1×1 arcmin² bins. The analysis is performed with the free-form code WSLAP+ (Diego et al. 2007; Sendra et al. 2014). This algorithm assumes the mass distribution can be decomposed as a superposition of 2D Gaussian functions. The amplitude of each Gaussian is optimized by minimizing the difference between the observed and predicted lensing variables (mean shear in the adopted 1×1 arcmin² bins for the WL portion of the data set, and positions of multiply imaged galaxies for the SL portion of the data set). The unknown position of the multiply imaged galaxies in the source plane is also optimized together with the distribution of mass. In the inner portion of the region constrained by the SL measurement, the light distribution of the most prominent member galaxies (approximately 70 for this cluster) is assumed to trace the mass on small scales with a given mass-to-light ratio, that is also derived as part of the optimization. The resulting convergence profile is shown in Fig. 9, where we show the solution obtained

when only SL data are used and when both SL and WL data are used. The profile in the inner region ($R \lesssim 0.44$ Mpc) contains the contribution from member galaxies. Even though member galaxies make only a small contribution to the profile of the convergence (except in the central peak), these galaxies are relevant for accurately predicting the positions of the multiply-imaged lensed galaxies and they also act as a regularization mechanism for the solution to converge (see Diego et al. 2007; Sendra et al. 2014). The SL+WL profile differs significantly from the SL profile at radii beyond the region constrained by SL. This is a consequence of the SL-only solution not being constrained in this regime, where WSLAP+ solutions are known to be biased low (this is a memory effect of the initial guess as described in Diego et al. 2007). In contrast, the SL+WL combination recovers a solution consistent with the expected Navarro-Frenk-White (NFW) profile (Navarro et al. 1996) of massive clusters at large radii.

8. *Euclid*'s preview of high-redshift galaxy clusters detected in the field of A2764

In order to illustrate the unique capabilities of *Euclid* to detect high-redshift ($z > 0.6$) galaxy clusters, we have searched for concentrations of galaxies in the VIS and NISP images in the direction of already known high-redshift clusters. To do so, we have focused on the increasingly large numbers of clusters that are blindly detected with surveys in the millimeter spectral range via the Sunyaev-Zeldovich effect (SZ; Sunyaev & Zeldovich 1972b,a, 1980; Planck Collaboration: Ade et al. 2016; Planck Collaboration: Adam et al. 2016; Bleem et al. 2024) and in the X-ray range (e.g. Bulbul et al. 2024). More specifically, we used ground-based telescopes such as the Atacama Cosmology Telescope (ACT, e.g. Hilton et al. 2021) and the South Pole Telescope (SPT, e.g. Bleem et al. 2015) which allow us to detect clusters out to $z \sim 1.9$. We also make use of the eROSITA-DE DR1 point and extended sources (Merloni et al. 2024). As shown in Bulbul et al. (2024), the first release of eROSITA data is already able to identify groups and clusters at high redshift, out to $z \sim 1.3$.

The ERO field of A2764 is located in both the ACT and SPT footprints. We searched for the presence of clusters from these two surveys in the ERO observation. We found that the A2764 ERO field contains a few background galaxy clusters previously detected in SZ with masses $M_{500} \sim 2 \times 10^{14} M_{\odot}$, located in the redshift range $z \sim 0.6-0.7$, which is higher than the target cluster. Three of them are ACT clusters (Hilton et al. 2021, DR5): (1) ACT-CLJ0022.5-4843 is an ACT-only cluster at $z_{\text{phot}} = 0.855 \pm 0.015$ with an estimated mass of $M_{500} = 1.71^{+0.37}_{-0.30} \times 10^{14} M_{\odot}$; (2) ACT-CLJ0021.5-4902 at $z_{\text{spec}} = 0.688$ and $M_{500} = 2.35^{+0.45}_{-0.38} \times 10^{14} M_{\odot}$ is also identified as SPT cluster SPT-CLJ0021.5-4902 (see Bleem et al. 2015; Everett et al. 2020) and it is associated with an eROSITA point-source (see below); and (3) ACT-CLJ0023.7-4923 with photometric redshift $z_{\text{phot}} = 0.709 \pm 0.021$ and mass $M_{500} = 1.88^{+0.41}_{-0.33} \times 10^{14} M_{\odot}$. These three clusters are clearly identified in the VIS and NISP images as concentrations of galaxies. Moreover, we note that both ACT-CLJ0023.7-4923 and ACT-CLJ0021.5-4902 show strong gravitational lensing features observed both in the VIS and NISP images (see Figs. 10 and 11). To our knowledge, only the strong lensing in ACT-CLJ0023.7-4923 was previously known (Diehl et al. 2017), thus the strong lensing observed in ACT-CLJ0021.5-4902 is a first detection for this cluster. Finally, a faint SPT source, SPT-SJ002330-4947.3, detected at 150 GHz (Everett et al. 2020) is located near the noisy edge of the A2764




Fig. 8. WL convergence κ reconstruction of the A2390 field. The left panel shows the reconstructed E-mode signal, with the cluster detected in the central region with high significance. The right panel shows the B mode, which does not include a cosmological signal to first order, but provides an illustration for the noise level of the reconstruction. The green contours start at $\kappa = 0.03$ and are in steps of $\Delta\kappa = 0.03$. Both reconstructions assume a zero mean convergence when averaged over the full reconstructed sky area.




Fig. 9. Mass profile of A2390 from an SL-only analysis and a joint SL+WL analysis. The two regions that include members+Gaussians ($R \leq 0.44$ Mpc) in the lens model or only Gaussian functions are indicated. We also show the range of distances covered by the SL constraints (from the brightest cluster galaxy). For the SL+WL combination, we show a range of models that are consistent with the data (≈ 1 standard deviation). The addition of WL data from *Euclid* is key to properly constrain the mass of the cluster out to the virial radius (corresponding to R_{200} or radius at which the density reaches 200 times the mean density of the Universe) with a vertical orange bar, where the width is the uncertainty in the virial radius.

Euclid ERO field. The quality of the images does not allow us to identify the possible visible or NIR counterparts.

More than 40 point and extended sources from the eROSITA DR1 catalog (Merloni et al. 2024) are found in the A2764 ERO field. Obviously, the cluster A2764 itself is among the eROSITA DR1 sources in the field. It appears as the two extended sources 1eRASSJ002035.1–491245 and 1eRASSJ002032.1–491432. The two ACT galaxy clusters (Hilton et al. 2021) are matched with eROSITA DR1 point-sources: ACT-CLJ0021.5–4902 at $z_{\text{spec}} = 0.688$ is associated with 1eRASSJ002132.6–490216 (IDCluster=758, EXT=0) and ACT-CLJ0023.7–4923 at $z_{\text{phot}} = 0.709 \pm 0.021$ is associated with 1eRASSJ002345.0–492354 (ID-

Cluster=273 EXT=0). Only one source labelled as extended in the eROSITA DR1 catalog is found in the A2764 ERO field, and would qualify as a galaxy cluster: 1eRASSJ002223.6–492746 (with IDCluster=183, EXT=19.3 and EXT_LIKE=3.07). The VIS and NISP images show concentrations of galaxies at the associated location, and the distribution of photometric redshifts indicates a peak at $z_{\text{phot}} \sim 0.8$. We also checked for counterparts in the Massive and Distant Clusters of WISE Survey (MaD-CoWs) catalogs (Gonzalez et al. 2019) and didn't find any match in the ERO A2764 field.

Another remarkable eROSITA DR1 point source is 1eRASSJ002212.5–492811 (IDCluster=47 EXT=0). It is less than $2'$ away from the aforementioned source 1eRASSJ002223.6–492746 and about 40 VIS and NISP galaxies are located in its vicinity, half a dozen with a photometric redshift peaking at $z_{\text{phot}} \sim 0.8$ and about 30 at $z_{\text{phot}} \sim 1.4$. Although the association of *Euclid* galaxies and this eROSITA source is not yet confirmed, it illustrates the power of the *Euclid* survey in combination with other wavelength datasets to efficiently provide robust galaxy cluster candidates.

This simple search for optical and near-IR counterparts to already known SZ clusters at $z > 0.6$ with $M_{500} > 10^{14} M_{\odot}$ in this ERO program data demonstrates that *Euclid* VIS and NISP instruments are particularly efficient at detecting clusters at $z > 0.6$. Matching with eROSITA extended and point-sources also opens the way for the detection of new clusters at potentially even higher redshifts. Further work is needed to confirm associations between optical/NIR *Euclid* galaxies with millimeter or X-ray-detected clusters. We have also yet to run dedicated cluster-finding algorithms to find galaxy clusters and proto-clusters in the *Euclid* data. Nevertheless, this first analysis demonstrates that *Euclid* data will be a treasure trove for cluster studies.

9. Intracluster light

Intracluster light (ICL) is diffuse light seen in galaxy groups and clusters that comes from stars that are bound to the gravitational potential well of the cluster rather than any particular galaxy (see Contini 2021; Montes 2022, for reviews). It is a prominent





Fig. 10. $30'' \times 30''$ cutout centered on the galaxy cluster ACT-CLJ0021.5–4902 at $z_{\text{spec}} = 0.688$, associated with the eROSITA DR1 source 1eRASSJ002132.6–490216, as seen in VIS I_E (top panel) and NISP H_E (bottom panel). We can clearly identify a strong lensing arc that was previously unknown.

Fig. 11. $30'' \times 30''$ cutout centered on the galaxy cluster ACT-CLJ0023.7–4923 with $z_{\text{phot}} = 0.709 \pm 0.021$, associated with the eROSITA DR1 source 1eRASSJ002345.0–492354, as seen in VIS I_E (top panel) and NISP H_E (bottom panel). Multiple lensing features, which are already known, are visible in the image. The star on the right of the image is *Gaia* 4973928409193111936 of $G_{\text{RP}} = 14$ mag.

feature observed in clusters (e.g. Feldmeier et al. 2004; Kluge et al. 2020; Golden-Marx et al. 2023), and the origin and assembly history of the ICL serves as a tool to characterize the assembly history of the brightest central galaxy and the satellite galaxy population (e.g., Murante et al. 2007). In addition, the ICL can be used as a luminous tracer of the dark matter distribution within the cluster (Montes & Trujillo 2019; Alonso Asensio et al. 2020; Deason et al. 2021; Gonzalez et al. 2021; Yoo et al. 2022; Contreras-Santos et al. 2024).

Although the ICL appears to be a powerful tool for studying galaxy clusters and cosmology, our understanding of this component has been hampered by observational limitations. For example, ground-based observations have been challenging due to the extreme faintness of this emission. HST has provided ultra-deep images of galaxy clusters, but it has a clear limitation: its small field of view. The high resolution, space-based images of *Euclid* can characterize the ICL beyond the typical HST limit of 100–200 kpc and will help improve our understanding of the formation of this diffuse light and the assembly of clusters. In this




Fig. 12. H_E -band image showing the extent of ICL in A2390. Stars in the field have been subtracted using the PSF model of Cuillandre et al. (2024) and the diffraction spikes have been masked by hand. The lognormal color scale is chosen to highlight the diffuse ICL in this cluster.

sense, *Euclid* can finally provide the wide field of view needed to probe the outer regions of clusters. Using observations from this ERO program, we will probe the ICL of A2390, A2764 and the other high redshift clusters detected in the field of A2764 out to distances of several hundred kpc.

Figure 12 shows the extent and distribution of the ICL surrounding A2390 in the H_E band. We used the outer PSF model of Cuillandre et al. (2024) to subtract the PSF contribution of several bright stars in the field. To highlight the low surface-brightness ICL, we masked the remaining diffraction spikes by hand and interpolated over the masked pixels using a Gaussian kernel with $\sigma = 0''.6$. The ICL stretches over approximately 1 Mpc, from southwest to northwest, following the distribution of the brightest members of the cluster. The 1σ image depth is

$\mu(H_E) = 28.88 \text{ mag arcsec}^{-2}$ measured over a $10'' \times 10''$ area; see Cuillandre et al. (2024) for details. At this depth, the main sources that limit the reliability of the ICL measurement are Galactic cirri and noise from the subtraction of the NIR persistence. The ERO data on the Perseus cluster reached a similar depth as these ERO data on A2390 and A2764 (Cuillandre et al. 2024). The ICL of the Perseus cluster was reliably measured out to a semi-major axis distance of 600 kpc reaching a 1σ depth of $\mu(I_E) = 29.4 \text{ mag arcsec}^{-2}$, $\mu(Y_E) = 28.4 \text{ mag arcsec}^{-2}$, and $\mu(J_E) = \mu(H_E) = 28.7 \text{ mag arcsec}^{-2}$ (Kluge et al. 2024) in similar cirri conditions and significantly more persistence (due to larger galaxies in the field of view). Hence, we expect to reach similar depths in these images.

10. Summary

The ERO program presents an opportunity to highlight the capabilities and performance of the *Euclid* spacecraft, offering a glimpse into the nominal survey observations. In this paper, we utilize data acquired from the ERO11 program, which focused on two lensing clusters, A2390 and A2764. The nature and depth of these data enable exploration of various scientific cases related to galaxy formation and evolution, weak and strong lensing phenomena, as well as the physics of galaxy clusters. We have produced a photometric catalog in I_E , Y_E , J_E , and H_E bands for a total of 501 994 sources using different aperture sizes. We also performed multi-band photometry based on VIS detection, which include supporting ground-based observations. These catalogs are made publicly available along with this paper.

The combination of NISP and VIS imaging is well-suited for identifying dropout sources that are visible only in the NISP images and remain undetected in the optical I_E filter. This is particularly relevant for detecting the Lyman break of high-redshift sources at $z > 6$, since the survey design naturally produces roughly 3 magnitudes breaks. Combining photometric criteria with SED-fitting with Eazy, we have identified a total of 30 LBG candidates. The compactness of these sources, which remain unresolved in the NISP images, presents a challenge in discriminating between galaxies and quasars. Furthermore, despite exhibiting an apparently pronounced Lyman break, certain rare brown dwarfs can display colors resembling those of our targeted sources. Nevertheless, considering that *Euclid* possesses the unique capability to efficiently identify such bright and rare sources on the sky at $z > 6$, it becomes clear that this will facilitate extensive spectroscopic follow-up campaigns from both ground- and space-based observatories, enabling the confirmation of their redshifts and the study of their physical characteristics.

These cluster fields also offer the possibility to perform a joint SL+WL analysis to model the mass distribution of these structures. We have combined an already existing SL model of A2390 with our own WL analysis to extend constraints to larger radius and better estimate the virial mass of the cluster. Using the WSLAP+ code, we show how WL improves the mass profile reconstruction of the cluster compared to SL only.

We have also illustrate *Euclid*'s potential for detecting high-redshift galaxy clusters by using these ERO data to search for optical and NIR counterparts of known clusters. In particular, we have identified three clusters in the A2764 field, which were previously detected by their X-ray emission (eROSITA) or by the SZ effect (ACT and SPT). Two of the three clusters, which are located at $z = 0.6$ – 0.9 show clear gravitational lensing features. This early work already demonstrates the ability of multi-wavelength efforts to discover high-redshift clusters and to calibrate, in the near-future, automatic identification algorithms.

Finally, cluster fields are ideal laboratories to study the ICL, which helps constrain the assembly history of cluster galaxies and to indirectly map the dark matter distribution. We illustrate the large extent of the ICL in A2390 visible in the *Euclid* images, which go far beyond the limited field of view of HST.

Overall, this paper underscores the large array of scientific studies that will be carried out in blank (but also lensed) fields with *Euclid*. The depth, angular resolution, and wide field of view are clearly a unique combination that will push many frontiers in the landscape of *Euclid*'s legacy science.

Acknowledgements. HA is supported by the French Centre National d'Etudes Spatial (CNES). The Cosmic Dawn Center (DAWN) is funded by the Danish National Research Foundation (DNRF140). This work has made use of the

CANDIDE Cluster at the *Institut d'Astrophysique de Paris* (IAP), made possible by grants from the PNCG and the region of Île de France through the program DIM-ACAV+, and the Cosmic Dawn Center and maintained by S. Rouberol. HM is supported by JSPS KAKENHI Grant Number JP23H00108, and JSPS Core-to-Core Program Grant Numbers JPJSCCA20200002 and JPJSCCA20210003. MJ is supported by the United Kingdom Research and Innovation (UKRI) Future Leaders Fellowship 'Using Cosmic Beasts to uncover the Nature of Dark Matter' (grants number MR/S017216/1 & MR/X006069/1). TS acknowledges support provided by the Austrian Research Promotion Agency (FFG) and the Federal Ministry of the Republic of Austria for Climate Action, Environment, Energy, Mobility, Innovation and Technology (BMK) via the Austrian Space Applications Programme with grant numbers 899537 and 900565, as well as the German Research Foundation (DFG) under grant 415537506, and the German Federal Ministry for Economic Affairs and Climate Action (BMWK) provided by DLR under projects no. 50QE2002 and 50QE2302.

This work has made use of the Early Release Observations (ERO) data from the *Euclid* mission of the European Space Agency (ESA), 2024, <https://doi.org/10.57780/esa-qmocz3>.

The *Euclid* Consortium acknowledges the European Space Agency and a number of agencies and institutes that have supported the development of *Euclid*, in particular the Agenzia Spaziale Italiana, the Austrian Forschungsförderungsgesellschaft, funded through BMK, the Belgian Science Policy, the Canadian *Euclid* Consortium, the Deutsches Zentrum für Luft- und Raumfahrt, the DTU Space and the Niels Bohr Institute in Denmark, the French Centre National d'Etudes Spatiales, the Fundação para a Ciência e a Tecnologia, the Hungarian Academy of Sciences, the Ministerio de Ciencia, Innovación y Universidades, the National Aeronautics and Space Administration, the National Astronomical Observatory of Japan, the Nederlandse Onderzoekschool Voor Astronomie, the Norwegian Space Agency, the Research Council of Finland, the Romanian Space Agency, the State Secretariat for Education, Research, and Innovation (SERI) at the Swiss Space Office (SSO), and the United Kingdom Space Agency. A complete and detailed list is available on the *Euclid* web site (<http://www.euclid-ec.org>).

This work made use of Astropy⁵: a community-developed core Python package and an ecosystem of tools and resources for astronomy (Astropy Collaboration et al. 2013, 2018, 2022) and Matplotlib (Hunter 2007). This research has made use of Aladin sky atlas, CDS, Strasbourg Astronomical Observatory, France (Bonnarel et al. 2000).

References

- Abbott, T. M. C., Adamów, M., Aguena, M., et al. 2021, *ApJS*, 255, 20
- Abell, G. O., Corwin, Harold G., J., & Olowin, R. P. 1989, *ApJS*, 70, 1
- Adams, N. J., Conselice, C. J., Ferreira, L., et al. 2023, *MNRAS*, 518, 4755
- Alonso Asensio, I., Dalla Vecchia, C., Bahé, Y. M., Barnes, D. J., & Kay, S. T. 2020, *MNRAS*, 494, 1859
- Applegate, D. E., von der Linden, A., Kelly, P. L., et al. 2014, *MNRAS*, 439, 48
- Astropy Collaboration, Price-Whelan, A. M., Lim, P. L., et al. 2022, *ApJ*, 935, 167
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, 156, 123
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33
- Atek, H., Chemerynska, I., Wang, B., et al. 2023, *MNRAS*, 524, 5486
- Barbary, K. 2016, *The Journal of Open Source Software*, 1, 58
- Bartelmann, M. & Schneider, P. 2001, *Phys. Rep.*, 340, 291
- Bertin, E. 2011, in *Astronomical Society of the Pacific Conference Series*, Vol. 442, *Astronomical Data Analysis Software and Systems XX*, ed. I. N. Evans, A. Accomazzi, D. J. Mink, & A. H. Rots, 435
- Bertin, E. & Arnouts, S. 1996, *A&AS*, 117, 393
- Bertin, E., Mellier, Y., Radovich, M., et al. 2002, in *Astronomical Society of the Pacific Conference Series*, Vol. 281, *Astronomical Data Analysis Software and Systems XI*, ed. D. A. Bohlender, D. Durand, & T. H. Handley, 228
- Bertin, E., Schefer, M., Apostolakis, N., et al. 2022, *SourceXtractor++: Extracts sources from astronomical images*, *Astrophysics Source Code Library*, record ascl:2212.018
- Bleem, L. E., Klein, M., Abbot, T. M. C., et al. 2024, *The Open Journal of Astrophysics*, 7, 13
- Bleem, L. E., Stalder, B., de Haan, T., et al. 2015, *ApJS*, 216, 27
- Bohringer, H., Tanaka, Y., Mushotzky, R. F., Ikebe, Y., & Hattori, M. 1998, *A&A*, 334, 789
- Bonnarel, F., Fernique, P., Bienaimé, O., et al. 2000, *A&AS*, 143, 33
- Bouwens, R. J., Illingworth, G. D., Oesch, P. A., et al. 2015, *ApJ*, 803, 34
- Bowler, R. A. A., Dunlop, J. S., McLure, R. J., & McLeod, D. J. 2017, *MNRAS*, 466, 3612

⁵ <http://www.astropy.org>

- Bowler, R. A. A., Jarvis, M. J., Dunlop, J. S., et al. 2020, *MNRAS*, 493, 2059
- Brammer, G., Strait, V., Matharu, J., & Momcheva, I. 2022, *grizli*, Zenodo
- Brammer, G. B., van Dokkum, P. G., & Coppi, P. 2008, *ApJ*, 686, 1503
- Bulbul, E., Liu, A., Kluge, M., et al. 2024, *arXiv e-prints*, [arXiv:2402.08452](https://arxiv.org/abs/2402.08452)
- Chemerynska, I., Atek, H., Furtak, L. J., et al. 2023, *arXiv e-prints*, [arXiv:2312.05030](https://arxiv.org/abs/2312.05030)
- Contini, E. 2021, *Galaxies*, 9, 60
- Contreras-Santos, A., Knebe, A., Cui, W., et al. 2024, *arXiv e-prints*, [arXiv:2401.08283](https://arxiv.org/abs/2401.08283)
- Cowie, L. L., Barger, A. J., Bauer, F. E., et al. 2022, *ApJ*, 939, 5
- Cuillandre, J.-C., Bertin, E., Bolzonella, M., et al. 2024, *A&A*, this issue
- Dark Energy Survey Collaboration, Abbott, T., Abdalla, F. B., et al. 2016, *MNRAS*, 460, 1270
- Deason, A. J., Oman, K. A., Fattahi, A., et al. 2021, *MNRAS*, 500, 4181
- Dey, A., Schlegel, D. J., Lang, D., et al. 2019, *AJ*, 157, 168
- Diego, J. M., Tegmark, M., Protopapas, P., & Sandvik, H. B. 2007, *MNRAS*, 375, 958
- Diehl, H. T., Buckley-Geer, E. J., Lindgren, K. A., et al. 2017, *ApJS*, 232, 15
- Donnan, C. T., McLure, R. J., Dunlop, J. S., et al. 2024, *arXiv e-prints*, [arXiv:2403.03171](https://arxiv.org/abs/2403.03171)
- Erben, T., Van Waerbeke, L., Bertin, E., Mellier, Y., & Schneider, P. 2001, *A&A*, 366, 717
- Euclid Collaboration: Barnett, R., Warren, S. J., Mortlock, D. J., et al. 2019, *A&A*, 631, A85
- Euclid Collaboration: Bretonnière, H., Kuchner, U., Huertas-Company, M., et al. 2023, *A&A*, 671, A102
- Euclid Collaboration: Cropper, M., Al Bahlawan, A., Amiaux, J., et al. 2024, *A&A*, this issue
- Euclid Collaboration: Jahnke, K., Gillard, W., Schirmer, M., et al. 2024, *A&A*, this issue
- Euclid Collaboration: Martinet, N., Schrabback, T., Hoekstra, H., et al. 2019, *A&A*, 627, A59
- Euclid Collaboration: Mellier, Y., Abdurro'uf, Acevedo Barroso, J., Achúcarro, A., et al. 2024, *A&A*, this issue
- Euclid Collaboration: Merlin, E., Castellano, M., Bretonnière, H., et al. 2023, *A&A*, 671, A101
- Euclid Collaboration: Scaramella, R., Amiaux, J., Mellier, Y., et al. 2022, *A&A*, 662, A112
- Euclid Collaboration: van Mierlo, S. E., Caputi, K. I., Ashby, M., et al. 2022, *A&A*, 666, A200
- Everett, W. B., Zhang, L., Crawford, T. M., et al. 2020, *ApJ*, 900, 55
- Feix, M., Zhao, H., Fedeli, C., Pestaña, J. L. G., & Hoekstra, H. 2010, *Phys. Rev. D*, 82, 124003
- Feldmeier, J. J., Mihos, J. C., Morrison, H. L., et al. 2004, *ApJ*, 609, 617
- Finkelstein, S. L., Ryan, Jr., R. E., Papovich, C., et al. 2015, *ApJ*, 810, 71
- Fitzpatrick, E. L. & Massa, D. 2007, *ApJ*, 663, 320
- Flaugher, B. 2005, *International Journal of Modern Physics A*, 20, 3121
- Flaugher, B., Diehl, H. T., Honscheid, K., et al. 2015, *AJ*, 150, 150
- Golden-Marx, J. B., Zhang, Y., Ogando, R. L. C., et al. 2023, *MNRAS*, 521, 478
- Gonzalez, A. H., George, T., Connor, T., et al. 2021, *MNRAS*, 507, 963
- Gonzalez, A. H., Gettings, D. P., Brodwin, M., et al. 2019, *ApJS*, 240, 33
- Harikane, Y., Ouchi, M., Oguri, M., et al. 2023, *ApJS*, 265, 5
- Hernández-Martín, B., Schrabback, T., Hoekstra, H., et al. 2020, *A&A*, 640, A117
- Hilton, M., Sifón, C., Naess, S., et al. 2021, *ApJS*, 253, 3
- Hoekstra, H., Franx, M., Kuijken, K., & Squires, G. 1998, *ApJ*, 504, 636
- Hunter, J. D. 2007, *Computing in Science and Engineering*, 9, 90
- Kaiser, N., Squires, G., & Broadhurst, T. 1995, *ApJ*, 449, 460
- Katgert, P., Mazure, A., Perea, J., et al. 1996, *A&A*, 310, 8
- Kluge, M., Hatch, N., Montes, M., et al. 2024, *A&A*, this issue
- Kluge, M., Neureiter, B., Riffeser, A., et al. 2020, *ApJS*, 247, 43
- Koekemoer, A. M., Faber, S. M., Ferguson, H. C., et al. 2011, *ApJS*, 197, 36
- Kümmel, M., Álvarez-Ayllón, A., Bertin, E., et al. 2022, *arXiv e-prints*, [arXiv:2212.02428](https://arxiv.org/abs/2212.02428)
- Le Borgne, J. F., Mathez, G., Mellier, Y., et al. 1991, *A&AS*, 88, 133
- Liu, Q., Yee, H. K. C., Drissen, L., et al. 2021, *ApJ*, 908, 228
- Luppino, G. A. & Kaiser, N. 1997, *ApJ*, 475, 20
- Mandelbaum, R., Rowe, B., Armstrong, R., et al. 2015, *MNRAS*, 450, 2963
- McInnes, R. N., Menanteau, F., Heavens, A. F., et al. 2009, *MNRAS*, 399, L84
- Merloni, A., Lamer, G., Liu, T., et al. 2024, *A&A*, 682, A34
- Miyazaki, S., Komiyama, Y., Sekiguchi, M., et al. 2002, *PASJ*, 54, 833
- Montes, M. 2022, *Nature Astronomy*, 6, 308
- Montes, M. & Trujillo, I. 2019, *MNRAS*, 482, 2838
- Murante, G., Giovalli, M., Gerhard, O., et al. 2007, *MNRAS*, 377, 2
- Navarro, J. F., Frenk, C. S., & White, S. D. M. 1996, *ApJ*, 462, 563
- Okabe, N. & Smith, G. P. 2016, *MNRAS*, 461, 3794
- Olmstead, A., Rigby, J. R., Swinbank, M., & Veilleux, S. 2014, *AJ*, 148, 65
- Ota, K., Richard, J., Iye, M., et al. 2012, *MNRAS*, 423, 2829
- Ouchi, M., Shimasaku, K., Okamura, S., et al. 2004, *ApJ*, 611, 685
- Pello, R., Le Borgne, J.-F., Soucail, G., Mellier, Y., & Sanahuja, B. 1991, *ApJ*, 366, 405
- Planck Collaboration: Adam, R., Aghanim, N., Ashdown, M., et al. 2016, *A&A*, 596, A108
- Planck Collaboration: Ade, P. A. R., Aghanim, N., Arnaud, M., et al. 2016, *A&A*, 594, A13
- Richard, J., Claeysens, A., Lagattuta, D., et al. 2021, *A&A*, 646, A83
- Schneider, P. & Er, X. 2008, *A&A*, 485, 363
- Schrabback, T., Applegate, D., Dietrich, J. P., et al. 2018, *MNRAS*, 474, 2635
- Schrabback, T., Bocquet, S., Sommer, M., et al. 2021, *MNRAS*, 505, 3923
- Sendra, I., Diego, J. M., Broadhurst, T., & Lazkoz, R. 2014, *MNRAS*, 437, 2642
- Sereno, M. & Ettori, S. 2017, *MNRAS*, 468, 3322
- Simon, P., Taylor, A. N., & Hartlap, J. 2009, *MNRAS*, 399, 48
- Sohn, J., Fabricant, D. G., Geller, M. J., Hwang, H. S., & Diaferio, A. 2020, *ApJ*, 902, 17
- Sunyaev, R. A. & Zeldovich, I. B. 1980, *ARA&A*, 18, 537
- Sunyaev, R. A. & Zeldovich, Y. B. 1972a, *A&A*, 20, 189
- Sunyaev, R. A. & Zeldovich, Y. B. 1972b, *Comments on Astrophysics and Space Physics*, 4, 173
- von der Linden, A., Allen, M. T., Applegate, D. E., et al. 2014, *MNRAS*, 439, 2
- Weaver, J., Taamoli, S., McPartland, C., et al. 2024, *A&A*, this issue
- Yagi, M., Kashikawa, N., Sekiguchi, M., et al. 2002, *AJ*, 123, 66
- Yoo, J., Ko, J., Sabiu, C. G., et al. 2022, *ApJS*, 261, 28
- Zohren, H., Schrabback, T., Bocquet, S., et al. 2022, *A&A*, 668, A18

-
- ¹ Institut d'Astrophysique de Paris, UMR 7095, CNRS, and Sorbonne Université, 98 bis boulevard Arago, 75014 Paris, France
 - ² Aix-Marseille Université, CNRS, CNES, LAM, Marseille, France
 - ³ Department of Astronomy, University of Massachusetts, Amherst, MA 01003, USA
 - ⁴ Instituto de Física de Cantabria, Edificio Juan Jordá, Avenida de los Castros, 39005 Santander, Spain
 - ⁵ Universität Innsbruck, Institut für Astro- und Teilchenphysik, Technikerstr. 25/8, 6020 Innsbruck, Austria
 - ⁶ School of Physics and Astronomy, University of Nottingham, University Park, Nottingham NG7 2RD, UK
 - ⁷ Université Paris-Saclay, CNRS, Institut d'astrophysique spatiale, 91405, Orsay, France
 - ⁸ Department of Astronomy, University of Geneva, ch. d'Ecogia 16, 1290 Versoix, Switzerland
 - ⁹ Physics and Astronomy Department, University of California, 900 University Ave., Riverside, CA 92521, USA
 - ¹⁰ Institute for Astronomy, University of Edinburgh, Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, UK
 - ¹¹ Instituto de Astrofísica de Andalucía, CSIC, Glorieta de la Astronomía, 18080, Granada, Spain
 - ¹² Observatorio Nacional, Rua General Jose Cristino, 77-Bairro Imperial de Sao Cristovao, Rio de Janeiro, 20921-400, Brazil
 - ¹³ Université Paris-Saclay, Université Paris Cité, CEA, CNRS, AIM, 91191, Gif-sur-Yvette, France
 - ¹⁴ Max-Planck-Institut für Astronomie, Königstuhl 17, 69117 Heidelberg, Germany
 - ¹⁵ INFN-Osservatorio di Astrofisica e Scienza dello Spazio di Bologna, Via Piero Gobetti 93/3, 40129 Bologna, Italy
 - ¹⁶ Jodrell Bank Centre for Astrophysics, Department of Physics and Astronomy, University of Manchester, Oxford Road, Manchester M13 9PL, UK
 - ¹⁷ The University of Texas at Austin, Austin, TX, 78712, USA
 - ¹⁸ Department of Physics, Centre for Extragalactic Astronomy, Durham University, South Road, DH1 3LE, UK
 - ¹⁹ Department of Physics, Institute for Computational Cosmology, Durham University, South Road, DH1 3LE, UK
 - ²⁰ Astrophysics Research Centre, University of KwaZulu-Natal, Westville Campus, Durban 4041, South Africa
 - ²¹ School of Mathematics, Statistics & Computer Science, University of KwaZulu-Natal, Westville Campus, Durban 4041, South Africa
 - ²² STAR Institute, Quartier Agora - Allée du six Août, 19c B-4000 Liège, Belgium
 - ²³ Centre de Recherche Astrophysique de Lyon, UMR5574, CNRS, Université Claude Bernard Lyon 1, ENS de Lyon, 69230, Saint-Genis-Laval, France

- ²⁴ School of Physical Sciences, The Open University, Milton Keynes, MK7 6AA, UK
- ²⁵ Cosmic Dawn Center (DAWN)
- ²⁶ Niels Bohr Institute, University of Copenhagen, Jagtvej 128, 2200 Copenhagen, Denmark
- ²⁷ Dipartimento di Fisica "Aldo Pontremoli", Università degli Studi di Milano, Via Celoria 16, 20133 Milano, Italy
- ²⁸ INAF-Osservatorio Astronomico di Trieste, Via G. B. Tiepolo 11, 34143 Trieste, Italy
- ²⁹ IFPU, Institute for Fundamental Physics of the Universe, via Beirut 2, 34151 Trieste, Italy
- ³⁰ INAF-Osservatorio Astronomico di Roma, Via Frascati 33, 00078 Monteporzio Catone, Italy
- ³¹ INAF-IASF Milano, Via Alfonso Corti 12, 20133 Milano, Italy
- ³² Astrophysics Group, Blackett Laboratory, Imperial College London, London SW7 2AZ, UK
- ³³ Institute of Physics, Laboratory of Astrophysics, Ecole Polytechnique Fédérale de Lausanne (EPFL), Observatoire de Sauverny, 1290 Versoix, Switzerland
- ³⁴ Institut d'Astrophysique de Paris, 98bis Boulevard Arago, 75014, Paris, France
- ³⁵ Cosmic Dawn Center (DAWN), Denmark
- ³⁶ INFN-Sezione di Bologna, Viale Berti Pichat 6/2, 40127 Bologna, Italy
- ³⁷ Kobayashi-Maskawa Institute for the Origin of Particles and the Universe, Nagoya University, Chikusa-ku, Nagoya, 464-8602, Japan
- ³⁸ Institute for Advanced Research, Nagoya University, Chikusa-ku, Nagoya, 464-8601, Japan
- ³⁹ Kavli Institute for the Physics and Mathematics of the Universe (WPI), University of Tokyo, Kashiwa, Chiba 277-8583, Japan
- ⁴⁰ Instituto de Astrofísica de Canarias, Calle Vía Láctea s/n, 38204, San Cristóbal de La Laguna, Tenerife, Spain
- ⁴¹ Departamento de Astrofísica, Universidad de La Laguna, 38206, La Laguna, Tenerife, Spain
- ⁴² Department of Mathematics, Imperial College London, London SW7 2AZ, UK
- ⁴³ Physics Program, Graduate School of Advanced Science and Engineering, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-8526, Japan
- ⁴⁴ Hiroshima Astrophysical Science Center, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-8526, Japan
- ⁴⁵ Core Research for Energetic Universe, Hiroshima University, 1-3-1, Kagamiyama, Higashi-Hiroshima, Hiroshima 739-8526, Japan
- ⁴⁶ Dipartimento di Fisica e Scienze della Terra, Università degli Studi di Ferrara, Via Giuseppe Saragat 1, 44122 Ferrara, Italy
- ⁴⁷ Universitäts-Sternwarte München, Fakultät für Physik, Ludwig-Maximilians-Universität München, Scheinerstrasse 1, 81679 München, Germany
- ⁴⁸ Max Planck Institute for Extraterrestrial Physics, Giessenbachstr. 1, 85748 Garching, Germany
- ⁴⁹ European Space Agency/ESTEC, Keplerlaan 1, 2201 AZ Noordwijk, The Netherlands
- ⁵⁰ Observatoire Astronomique de Strasbourg (ObAS), Université de Strasbourg - CNRS, UMR 7550, Strasbourg, France
- ⁵¹ Kapteyn Astronomical Institute, University of Groningen, PO Box 800, 9700 AV Groningen, The Netherlands
- ⁵² Department of Physics, Université de Montréal, 2900 Edouard Montpetit Blvd, Montréal, Québec H3T 1J4, Canada
- ⁵³ Aurora Technology for European Space Agency (ESA), Camino bajo del Castillo, s/n, Urbanización Villafranca del Castillo, Villanueva de la Cañada, 28692 Madrid, Spain
- ⁵⁴ ESAC/ESA, Camino Bajo del Castillo, s/n., Urb. Villafranca del Castillo, 28692 Villanueva de la Cañada, Madrid, Spain
- ⁵⁵ School of Mathematics and Physics, University of Surrey, Guildford, Surrey, GU2 7XH, UK
- ⁵⁶ INAF-Osservatorio Astronomico di Brera, Via Brera 28, 20122 Milano, Italy
- ⁵⁷ Dipartimento di Fisica e Astronomia, Università di Bologna, Via Gobetti 93/2, 40129 Bologna, Italy
- ⁵⁸ INAF-Osservatorio Astronomico di Padova, Via dell'Osservatorio 5, 35122 Padova, Italy
- ⁵⁹ Centre National d'Etudes Spatiales – Centre spatial de Toulouse, 18 avenue Edouard Belin, 31401 Toulouse Cedex 9, France
- ⁶⁰ INAF-Osservatorio Astrofisico di Torino, Via Osservatorio 20, 10025 Pino Torinese (TO), Italy
- ⁶¹ Dipartimento di Fisica, Università di Genova, Via Dodecaneso 33, 16146, Genova, Italy
- ⁶² INFN-Sezione di Genova, Via Dodecaneso 33, 16146, Genova, Italy
- ⁶³ Department of Physics "E. Pancini", University Federico II, Via Cinthia 6, 80126, Napoli, Italy
- ⁶⁴ INAF-Osservatorio Astronomico di Capodimonte, Via Moiarriello 16, 80131 Napoli, Italy
- ⁶⁵ INFN section of Naples, Via Cinthia 6, 80126, Napoli, Italy
- ⁶⁶ Instituto de Astrofísica e Ciências do Espaço, Universidade do Porto, CAUP, Rua das Estrelas, PT4150-762 Porto, Portugal
- ⁶⁷ Faculdade de Ciências da Universidade do Porto, Rua do Campo de Alegre, 4150-007 Porto, Portugal
- ⁶⁸ Dipartimento di Fisica, Università degli Studi di Torino, Via P. Giuria 1, 10125 Torino, Italy
- ⁶⁹ INFN-Sezione di Torino, Via P. Giuria 1, 10125 Torino, Italy
- ⁷⁰ Mullard Space Science Laboratory, University College London, Holmbury St Mary, Dorking, Surrey RH5 6NT, UK
- ⁷¹ INFN-Sezione di Roma, Piazzale Aldo Moro, 2 - c/o Dipartimento di Fisica, Edificio G. Marconi, 00185 Roma, Italy
- ⁷² Centro de Investigaciones Energéticas, Medioambientales y Tecnológicas (CIEMAT), Avenida Complutense 40, 28040 Madrid, Spain
- ⁷³ Port d'Informació Científica, Campus UAB, C. Albareda s/n, 08193 Bellaterra (Barcelona), Spain
- ⁷⁴ Institute for Theoretical Particle Physics and Cosmology (TTK), RWTH Aachen University, 52056 Aachen, Germany
- ⁷⁵ Institute of Space Sciences (ICE, CSIC), Campus UAB, Carrer de Can Magrans, s/n, 08193 Barcelona, Spain
- ⁷⁶ Institut d'Estudis Espacials de Catalunya (IEEC), Edifici RDIT, Campus UPC, 08860 Castelldefels, Barcelona, Spain
- ⁷⁷ Dipartimento di Fisica e Astronomia "Augusto Righi" - Alma Mater Studiorum Università di Bologna, Viale Berti Pichat 6/2, 40127 Bologna, Italy
- ⁷⁸ European Space Agency/ESRIN, Largo Galileo Galilei 1, 00044 Frascati, Roma, Italy
- ⁷⁹ Université Claude Bernard Lyon 1, CNRS/IN2P3, IP2I Lyon, UMR 5822, Villeurbanne, F-69100, France
- ⁸⁰ UCB Lyon 1, CNRS/IN2P3, IUF, IP2I Lyon, 4 rue Enrico Fermi, 69622 Villeurbanne, France
- ⁸¹ Departamento de Física, Faculdade de Ciências, Universidade de Lisboa, Edifício C8, Campo Grande, PT1749-016 Lisboa, Portugal
- ⁸² Instituto de Astrofísica e Ciências do Espaço, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal
- ⁸³ INAF-Istituto di Astrofisica e Planetologia Spaziali, via del Fosso del Cavaliere, 100, 00100 Roma, Italy
- ⁸⁴ INFN-Padova, Via Marzolo 8, 35131 Padova, Italy
- ⁸⁵ School of Physics, HH Wills Physics Laboratory, University of Bristol, Tyndall Avenue, Bristol, BS8 1TL, UK
- ⁸⁶ Aix-Marseille Université, CNRS/IN2P3, CPPM, Marseille, France
- ⁸⁷ Istituto Nazionale di Fisica Nucleare, Sezione di Bologna, Via Irnerio 46, 40126 Bologna, Italy
- ⁸⁸ FRACTAL S.L.N.E., calle Tulipán 2, Portal 13 1A, 28231, Las Rozas de Madrid, Spain
- ⁸⁹ Institute of Theoretical Astrophysics, University of Oslo, P.O. Box 1029 Blindern, 0315 Oslo, Norway
- ⁹⁰ Leiden Observatory, Leiden University, Einsteinweg 55, 2333 CC Leiden, The Netherlands
- ⁹¹ Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA, 91109, USA
- ⁹² Department of Physics, Lancaster University, Lancaster, LA1 4YB, UK

- ⁹³ Felix Hormuth Engineering, Goethestr. 17, 69181 Leimen, Germany
- ⁹⁴ Technical University of Denmark, Elektrovej 327, 2800 Kgs. Lyngby, Denmark
- ⁹⁵ NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA
- ⁹⁶ Department of Physics and Helsinki Institute of Physics, Gustaf Hällströmin katu 2, 00014 University of Helsinki, Finland
- ⁹⁷ Université de Genève, Département de Physique Théorique and Centre for Astroparticle Physics, 24 quai Ernest-Ansermet, CH-1211 Genève 4, Switzerland
- ⁹⁸ Department of Physics, P.O. Box 64, 00014 University of Helsinki, Finland
- ⁹⁹ Helsinki Institute of Physics, Gustaf Hällströmin katu 2, University of Helsinki, Helsinki, Finland
- ¹⁰⁰ Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, UK
- ¹⁰¹ NOVA optical infrared instrumentation group at ASTRON, Oude Hoogeveensedijk 4, 7991PD, Dwingeloo, The Netherlands
- ¹⁰² INFN-Sezione di Milano, Via Celoria 16, 20133 Milano, Italy
- ¹⁰³ Universität Bonn, Argelander-Institut für Astronomie, Auf dem Hügel 71, 53121 Bonn, Germany
- ¹⁰⁴ Dipartimento di Fisica e Astronomia "Augusto Righi" - Alma Mater Studiorum Università di Bologna, via Piero Gobetti 93/2, 40129 Bologna, Italy
- ¹⁰⁵ Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, Laboratoire Lagrange, Bd de l'Observatoire, CS 34229, 06304 Nice cedex 4, France
- ¹⁰⁶ Université Paris Cité, CNRS, Astroparticule et Cosmologie, 75013 Paris, France
- ¹⁰⁷ Institut de Física d'Altes Energies (IFAE), The Barcelona Institute of Science and Technology, Campus UAB, 08193 Bellaterra (Barcelona), Spain
- ¹⁰⁸ Department of Physics and Astronomy, University of Aarhus, Ny Munkegade 120, DK-8000 Aarhus C, Denmark
- ¹⁰⁹ Waterloo Centre for Astrophysics, University of Waterloo, Waterloo, Ontario N2L 3G1, Canada
- ¹¹⁰ Department of Physics and Astronomy, University of Waterloo, Waterloo, Ontario N2L 3G1, Canada
- ¹¹¹ Perimeter Institute for Theoretical Physics, Waterloo, Ontario N2L 2Y5, Canada
- ¹¹² Space Science Data Center, Italian Space Agency, via del Politecnico snc, 00133 Roma, Italy
- ¹¹³ Institute of Space Science, Str. Atomistilor, nr. 409 Măgurele, Ilfov, 077125, Romania
- ¹¹⁴ Dipartimento di Fisica e Astronomia "G. Galilei", Università di Padova, Via Marzolo 8, 35131 Padova, Italy
- ¹¹⁵ Departamento de Física, FCFM, Universidad de Chile, Blanco Encalada 2008, Santiago, Chile
- ¹¹⁶ Satlantis, University Science Park, Sede Bld 48940, Leioa-Bilbao, Spain
- ¹¹⁷ Centre for Electronic Imaging, Open University, Walton Hall, Milton Keynes, MK7 6AA, UK
- ¹¹⁸ Infrared Processing and Analysis Center, California Institute of Technology, Pasadena, CA 91125, USA
- ¹¹⁹ Instituto de Astrofísica e Ciências do Espaço, Faculdade de Ciências, Universidade de Lisboa, Tapada da Ajuda, 1349-018 Lisboa, Portugal
- ¹²⁰ Universidad Politécnica de Cartagena, Departamento de Electrónica y Tecnología de Computadoras, Plaza del Hospital 1, 30202 Cartagena, Spain
- ¹²¹ Institut de Recherche en Astrophysique et Planétologie (IRAP), Université de Toulouse, CNRS, UPS, CNES, 14 Av. Edouard Belin, 31400 Toulouse, France
- ¹²² INFN-Bologna, Via Ippolito Nievo 46, 40126 Bologna, Italy
- ¹²³ Dipartimento di Fisica, Università degli studi di Genova, and INFN-Sezione di Genova, via Dodecaneso 33, 16146, Genova, Italy
- ¹²⁴ INFN, Sezione di Trieste, Via Valerio 2, 34127 Trieste TS, Italy
- ¹²⁵ SISSA, International School for Advanced Studies, Via Bonomea 265, 34136 Trieste TS, Italy
- ¹²⁶ INAF, Istituto di Radioastronomia, Via Piero Gobetti 101, 40129 Bologna, Italy
- ¹²⁷ Institut für Theoretische Physik, University of Heidelberg, Philosophenweg 16, 69120 Heidelberg, Germany
- ¹²⁸ Université St Joseph; Faculty of Sciences, Beirut, Lebanon
- ¹²⁹ Junia, EPA department, 41 Bd Vauban, 59800 Lille, France
- ¹³⁰ ICSC - Centro Nazionale di Ricerca in High Performance Computing, Big Data e Quantum Computing, Via Magnanelli 2, Bologna, Italy
- ¹³¹ Department of Physics and Astronomy, University of British Columbia, Vancouver, BC V6T 1Z1, Canada